

Gastroenterology State-wide Referral Criteria for Public Outpatient Services

This document is to be used as a guide for referrers and clinicians in NSW public outpatient services

The aim of SRC is to facilitate safe, timely and effective referral and prioritisation of patients requiring access to NSW public specialist outpatient services.

This document contains Gastroenterology SRC for the following presenting conditions:

- [Concern for colorectal cancer \(Rectal bleeding/Positive Faecal Occult Blood Test\) \(adult\)](#)
- [Iron deficiency \(adult\)](#)
- [Hepatocellular cancer \(suspected or known\)/Liver lesion \(adult\)](#)
- [Inflammatory/irritable bowel disease/syndrome \(suspected or known\) \(adult\)](#)
- [Liver dysfunction \(adult\)](#)
- [Cirrhosis \(suspected or known\) \(adult\)](#)
- [Upper gastrointestinal dysfunction \(adult\)](#)

Notes

- Gastroenterology SRC are applicable to NSW Local Health Districts and Specialty Health Networks with existing public gastroenterology and applicable allied health-led, nurse-led or medical/surgical-led outpatient services that manage the identified presenting conditions
- Gastroenterology SRC are applicable where the identified presenting conditions managed by gastroenterologists and surgeons are delivered in private practice as part of public-private hospital arrangements
- Gastroenterology SRC does not intend to change eligibility in terms of presenting conditions managed and referral sources accepted for NSW public gastroenterology and applicable allied health-led, nurse-led or medical/surgical-led outpatient services, however some NSW Local Health Districts and Specialty Health Networks may have different eligibility based on local context
- Gastroenterology SRC supports patients to be managed by the most appropriate member(s) of the multidisciplinary care team in the most appropriate setting based on their presenting condition
- Gastroenterology SRC sets thresholds for referral to NSW public gastroenterology and applicable allied health-led, nurse-led or medical/surgical-led outpatient services, and the expected clinical urgency category based on clinical need

Acknowledgements

NSW Health would like to extend its appreciation to all clinicians, administrators and consumers who contributed to the development of Gastroenterology SRC. NSW Health would also like to acknowledge the SRC developed by QLD Health, VIC Health, SA Health and WA Health which has been referenced to support the development of Gastroenterology SRC for NSW public outpatient services.

Significant contributors: Dr Miriam Levy (Clinical Lead), Dr Ian Norton, Dr Golo Ahlenstiel, Dr Steven Bollipo, Dr Tim Walker, Dr Peter Lim, Dr Ted Stoklosa, Dr Aravind Suppiah, Mark Costello, Kylie Bennetts, Kristine Dimagmaliw, Nicole Getreu, Dr Luke Morphet, Dr Daniel Ewald, Dr Ai-Vee Chua, Dr Anju Aggarwal, Dr Rowena Ivers, Dr Neil Merrett

Glossary

The structure for SRC in NSW is divided into the following five criterion.

In some cases, component(s) may not be applicable to each presenting condition. This will be denoted by 'Nil'.

Criterion	Description
Emergency	<ul style="list-style-type: none"> Clinical presentations or 'red flags' where referring health professionals should consider redirecting the patient to an Emergency Department or urgent care service, or seeking medical advice (e.g. phone on-call medical practitioner) These criteria should not be used by referring health professionals to refer to an NSW public specialist outpatient service
Out of scope / not routinely provided	<ul style="list-style-type: none"> Symptoms, conditions and/or presentations that would not routinely be provided by NSW public specialist outpatient services (i.e. could be optimally and safely managed in primary care) These criteria acknowledge and permit exceptions, where clinically appropriate
Access / prioritisation	<ul style="list-style-type: none"> Symptoms, conditions and/or presentations that advise referring health professionals, clinicians, patients and carers suitability for management by NSW public specialist outpatient services These criteria are classified by expected clinical urgency category and clinically recommended timeframes to be seen for a new outpatient appointment (i.e. Category 1: within 30 days, Category 2: within 90 days, Category 3: within 365 days) These criteria are only applicable where NSW public specialist outpatient services exist and manage the identified presenting condition
Referral requirements	<ul style="list-style-type: none"> Mandatory information that is to be supplied with referrals to NSW public specialist outpatient services for specific symptoms, conditions and/or presentations These criteria support with the determination of an appropriate clinical urgency category
Other content	<ul style="list-style-type: none"> Optional information that can be supplied with referrals to NSW public specialist outpatient services for specific symptoms, conditions and/or presentations These criteria may support with the determination of an appropriate clinical urgency category, however, are not required to continue referral processing

Gastroenterological emergencies

Note: this emergency criteria are not an exhaustive list of gastroenterological emergencies. Please refer to HealthPathways for more information.

Emergency Criteria
Abdominal mass with life-threatening symptoms where medical or surgical intervention is required
Complicated diarrhoea with severe bloody diarrhoea or if associated with profound dehydration or significantly systemically unwell
Melaena (dark stools) / haematemesis / vomiting in large volume or with haemodynamic compromise
Bowel obstruction
Acute pancreatitis
Suspected or known inflammatory bowel disease with symptoms suggestive of severe flare of known inflammatory bowel disease accompanied by fever, haemodynamic compromise, significant abdominal pain, suspected megacolon, perforation, bowel obstruction and/or abscess and unable to be controlled in the community
<p>Suspected or known inflammatory bowel disease with concern for severe or complicated colitis (perforation, toxic mega colon, abscess, bowel obstruction) indicated by:</p> <ul style="list-style-type: none"> • Fever • Tachycardia • Hypotension • Significant abdominal pain/peritonism • Abscess (abdominal or perianal) • Acute severe colitis: patients with > 6 bloody bowel motions per 24 hours plus at least one of the following: <ul style="list-style-type: none"> – Temperature > 37.8°C – Pulse rate > 90 bpm – Haemoglobin < 105 gm/L – Raised inflammatory markers (erythrocyte sedimentation rate (ESR) > 30 mm/hr or C-reactive protein (CRP) > 30 mg/L)
Gastrostomy feeding tube problems with blocked or displaced gastrostomy in patients with nil oral route for nutrition/hydration
Significant rectal bleeding with life-threatening symptoms where medical or surgical intervention is required
Acute, severe lower gastrointestinal tract bleeding in large volume or with haemodynamic compromise

Iron deficiency where anaemia (Hb < Lower Limit of Normal) with haemodynamic compromise (e.g. angina, shortness of breath, congestive cardiac failure or pre-syncope)
Suspected or known cirrhosis with acute clinical decompensation event (e.g. encephalopathy, gastrointestinal bleeding or new onset ascites, especially if with pain, fever or other systemic symptoms)
Liver failure with acute or decompensated chronic liver disease emerging (bilirubin > 100, INR > 2.0) with clinical signs of decompensation such as encephalopathy, ascites or bleeding, or liver abscess
Acute liver failure (e.g. encephalopathy with liver test severe derangement, especially INR > 1.5 or bilirubin > 60)
Sepsis in a patient with cirrhosis
Suspected ascending cholangitis
Severe hepatic encephalopathy
Suspected spontaneous bacterial peritonitis
Dysphagia with obstruction from food or foreign body associated with inability to tolerate oral intake, especially if drooling or pooling of saliva

Out of scope / Not routinely provided

Presenting Condition	Out of scope / Not routinely provided
Concern for colorectal cancer (Rectal bleeding/Positive Faecal Occult Blood Test) (adult)	<ul style="list-style-type: none"> • Haemorrhoid surgery/banding (suggest surgical referral if medical therapy fails) • Anal fissure surgery (suggest surgical referral if medical therapy fails) • Persistent but unchanged symptoms previously investigated • Completed colonoscopy with adequate bowel preparation in the last 2 years for the same symptoms
Iron deficiency (adult)	<ul style="list-style-type: none"> • Anaemia or iron deficiency secondary to haematological, renal, dietary, physiological or gynaecological cause • Normochromic, normocytic anaemia with normal iron studies • Isolated low serum iron
Liver dysfunction (adult)	<ul style="list-style-type: none"> • Fatty liver with normal liver function tests and Fibrosis-4 (FIB 4) score below 1.3 (meaning no significant fibrosis evident)
Dyspepsia/Heartburn/Reflux/Helicobacter pylori infection (adult)	<ul style="list-style-type: none"> • Patients with controlled symptoms • Non-smoker and < 50 years old that ceases treatment and symptoms return • Belching • Halitosis

Presenting Conditions

Concern for colorectal cancer (Rectal bleeding/Positive Faecal Occult Blood Test) (adult)

Emergency	
<p><i>If any of the following are present or suspected, please refer the patient to the emergency department (via ambulance if necessary) or seek emergent medical advice via phone to on-call consultant/registrar.</i></p> <ul style="list-style-type: none"> Abdominal mass with life-threatening symptoms where medical or surgical intervention is required Acute, severe lower gastrointestinal tract bleeding in large volume or with haemodynamic compromise 	
Out of scope / not routinely provided	
<ul style="list-style-type: none"> Haemorrhoid surgery/banding (suggest colorectal surgical referral if medical therapy fails) Anal fissure surgery (suggest colorectal surgical referral if medical therapy fails) Persistent but unchanged symptoms previously investigated Completed colonoscopy with adequate bowel preparation in the last 2 years for the same symptoms 	
Access / prioritisation	
<p>Category 1 (clinically recommended to be seen within 30 calendar days)</p>	<ul style="list-style-type: none"> Positive Faecal Occult Blood Test (FOBT) in National Bowel Screening Program Age ≥ 40 years old and positive FOBT with any of the following symptoms: <ul style="list-style-type: none"> Rectal bleeding Change in bowel habit or any lower gastrointestinal symptoms Unexplained anaemia (Hb < Lower Limit of Normal) Age ≥ 50 years old and negative FOBT, but any of the following symptoms: <ul style="list-style-type: none"> Rectal bleeding Change in bowel habit or any lower gastrointestinal symptoms Unexplained anaemia (Hb < Lower Limit of Normal) Any age and $\geq 5\%$ unexplained weight loss in past 1 month or $\geq 10\%$ unexplained weight loss in past 6 months Any age and abnormal imaging concerning for colorectal or upper gastrointestinal cancer
<p>Category 2 (clinically recommended to be seen within 90 calendar days)</p>	<ul style="list-style-type: none"> Age ≤ 39 years old and positive FOBT with any of the following symptoms: <ul style="list-style-type: none"> Rectal bleeding

	<ul style="list-style-type: none"> ○ Change in bowel habit or any lower gastrointestinal symptoms ○ Unexplained anaemia (Hb < Lower Limit of Normal) ● Age 40-49 years old and negative FOBT, but any of the following symptoms: <ul style="list-style-type: none"> ○ Rectal bleeding ○ Change in bowel habit or any lower gastrointestinal symptoms ○ Unexplained anaemia (Hb < Lower Limit of Normal) ● Age ≤ 49 years old and change in bowel habit with elevated faecal calprotectin (> 100 mcg/g)
<p>Category 3 (clinically recommended to be seen within 365 calendar days)</p>	<ul style="list-style-type: none"> ● Age ≤ 39 years old, negative FOBT with any of the following symptoms: <ul style="list-style-type: none"> ○ Rectal bleeding ○ Change in bowel habit or any lower gastrointestinal symptoms ○ Unexplained anaemia (Hb < Lower Limit of Normal) ● Family history of colon cancer <ul style="list-style-type: none"> ○ Known or suspected hereditary colon cancer syndrome (age for colonoscopy depends on syndrome) ○ Suggestive of a moderate or high-risk colon cancer (see Cancer Council Guidelines for more information) ● Personal history of polyps or colon cancer and recommended for repeat colonoscopy (provide Charleston comorbidity index score for age > 75)
<p>Referral requirements</p>	
<ul style="list-style-type: none"> ● Patient and family history of gastrointestinal cancer, including age of diagnosis ● Documented duration and associated symptoms ● Weight loss (amount and timeframe) ● Full blood count ● Haematinics (iron studies, red blood cell count, folate, vitamin B12) 	
<p>Other content</p>	
<ul style="list-style-type: none"> ● Faecal Occult Blood Test (FOBT) result ● Electrolytes, urea and creatinine (EUC) ● Rectal examination result ● Relevant imaging results ● Current and previous colonoscopy results (including histology) ● If the patient identifies as Aboriginal and/or Torres Strait Islander ● If the patient is considered 'at risk' and/or among a vulnerable, disadvantaged or priority population 	

Iron deficiency (adult)

Emergency	
<p><i>If any of the following are present or suspected, please refer the patient to the emergency department (via ambulance if necessary) or seek emergent medical advice via phone to on-call consultant/registrar.</i></p> <ul style="list-style-type: none"> Iron deficiency where anaemia (Hb < Lower Limit of Normal) with haemodynamic compromise (e.g. angina, shortness of breath, congestive cardiac failure or pre-syncope) 	
Out of scope / not routinely provided	
<ul style="list-style-type: none"> Anaemia or iron deficiency secondary to haematological, renal, dietary, physiological or gynaecological cause Normochromic, normocytic anaemia with normal iron studies Isolated low serum iron 	
Access / prioritisation	
<p>Category 1 (clinically recommended to be seen within 30 calendar days)</p>	<ul style="list-style-type: none"> Age \geq 40 years old and positive Faecal Occult Blood Test (FOBT) Age \geq 50 years old and negative FOBT, but any of the following symptoms: <ul style="list-style-type: none"> Rectal bleeding Change in bowel habit or any lower gastrointestinal symptoms Unexplained iron deficiency with or without anaemia (Hb < Lower Limit of Normal) Any age and \geq 5% unexplained weight loss in past 1 month or \geq 10% unexplained weight loss in past 6 months
<p>Category 2 (clinically recommended to be seen within 90 calendar days)</p>	<ul style="list-style-type: none"> \geq 40 years old with all of the following present: <ul style="list-style-type: none"> Negative FOBT Negative coeliac serology Gastrointestinal symptoms Unexplained iron deficiency with or without anaemia < 39 years old and recurrent unexplained iron deficiency or positive FOBT (with or without gastrointestinal symptoms) Any age and serology suggestive of coeliac disease (new or uncontrolled)
<p>Category 3 (clinically recommended to be seen within 365 calendar days)</p>	<ul style="list-style-type: none"> < 39 years old, single episode unexplained iron deficiency and negative FOBT (with or without gastrointestinal symptoms)
Referral requirements	
<ul style="list-style-type: none"> Duration of symptoms Any previously received iron therapy (duration and timing) Menstrual history, familial haemoglobinopathies, blood donations Family history of gastrointestinal cancer or coeliac disease 	

- Dietary history, including red meat intake
- Weight loss (amount and timeframe)
- Non-steroidal anti-inflammatory use
- Full blood count
- Haematinics (iron studies, red blood cell count, folate, vitamin B12)
- Coeliac serology (total immunoglobulin A (IgA), tissue transglutaminase (tTG) +/- anti-endomysial antibody (EMA))

Other content

- Faecal Occult Blood Test (FOBT) result
- Previous endoscopy / histology reports
- Electrolytes, urea and creatinine (EUC)
- Liver function test result
- If the patient identifies as Aboriginal and/or Torres Strait Islander
- If the patient is considered 'at risk' and/or among a vulnerable, disadvantaged or priority population

Hepatocellular cancer (suspected or known)/Liver lesion (adult)

Emergency	
<p><i>If any of the following are present or suspected, please refer the patient to the emergency department (via ambulance if necessary) or seek emergent medical advice via phone to on-call consultant/registrar.</i></p> <ul style="list-style-type: none"> • Nil 	
Out of scope / not routinely provided	
<ul style="list-style-type: none"> • Nil 	
Access / prioritisation	
<p>Category 1 (clinically recommended to be seen within 30 calendar days)</p>	<ul style="list-style-type: none"> • Known chronic liver disease and any of the following are present: <ul style="list-style-type: none"> ○ New lesion (excluding simple cysts) ○ Lesion suggestive of hepatocellular cancer according to radiology report ○ Raised AFP • No underlying liver disease but radiology suggests lesion suspicious for malignancy
<p>Category 2 (clinically recommended to be seen within 90 calendar days)</p>	<ul style="list-style-type: none"> • Other liver lesions (unless imaging strongly suggests benign aetiology)
<p>Category 3 (clinically recommended to be seen within 365 calendar days)</p>	<ul style="list-style-type: none"> • Nil
Referral requirements	
<ul style="list-style-type: none"> • Liver imaging result (current and any previous) • Full blood count • Tumour markers (alpha fetoprotein (AFP), carbohydrate antigen 19-9 (Ca 19-9), carcinoembryonic antigen (CEA)) • Liver function test result • Renal function test result 	
Other content	
<ul style="list-style-type: none"> • History of underlying liver disease and/or cirrhosis • Hepatitis B serology (HBV sAg, sAb, core IgG) • Hepatitis C serology (HCV Ab) • If the patient identifies as Aboriginal and/or Torres Strait Islander • If the patient is considered 'at risk' and/or among a vulnerable, disadvantaged or priority population 	

Inflammatory/irritable bowel disease/syndrome (suspected or known) (adult)

Emergency

If any of the following are present or suspected, please refer the patient to the emergency department (via ambulance if necessary) or seek emergent medical advice via phone to on-call consultant/registrars.

- Suspected or known inflammatory bowel disease with symptoms suggestive of severe flare of known inflammatory bowel disease accompanied by fever, haemodynamic compromise, significant abdominal pain, suspected megacolon, perforation, bowel obstruction and/or abscess and unable to be controlled in the community
- Suspected or known inflammatory bowel disease with concern for severe or complicated colitis (perforation, toxic mega colon, abscess, bowel obstruction):
 - Fever
 - Tachycardia
 - Hypotension
 - Significant abdominal pain/peritonism
 - Abscess (abdominal or perianal)
 - Acute severe colitis: patients with > 6 bloody bowel motions per 24 hours plus at least one of the following:
 - Temperature > 37.8°C
 - Pulse rate > 90 bpm
 - Haemoglobin < 105 gm/L
 - Raised inflammatory markers (erythrocyte sedimentation rate (ESR) > 30 mm/hr or C-reactive protein (CRP) > 30 mg/L)

Out of scope / not routinely provided

- Nil

Access / prioritisation

Category 1

(clinically recommended to be seen within 30 calendar days)

- Known inflammatory bowel disease where infectious diarrhoea is excluded (by stool MC and S, oral contraceptive pill (OCP) and Clostridium difficile toxin) and any of the following are present:
 - Elevated faecal calprotectin (> 100 mcg/g)
 - New progressive, or warning gastrointestinal symptoms (e.g. abdominal pain, vomiting)
 - Critical factor:
 - Anaemia
 - Low albumin
 - Elevated erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP)
 - Iron deficiency
 - $\geq 5\%$ unexplained weight loss in past 1 month or $\geq 10\%$ unexplained weight loss in past 6 months
 - New, abnormal imaging
 - Perianal pain/fistulae suspected
- Suspected inflammatory bowel disease where chronic diarrhoea or other suspicious symptoms > 6 weeks with

	<p>elevated faecal calprotectin (> 100 mcg/g), and any of the following are present:</p> <ul style="list-style-type: none"> ○ New progressive, or warning gastrointestinal symptoms (e.g. abdominal pain, vomiting) ○ Laboratory critical factor: <ul style="list-style-type: none"> - Anaemia - Low albumin - Elevated ESR/CRP - Iron deficiency ○ ≥ 5% unexplained weight loss in past 1 month or ≥ 10% unexplained weight loss in past 6 months ○ Abnormal imaging suggesting inflammatory bowel disease
<p>Category 2 (clinically recommended to be seen within 90 calendar days)</p>	<ul style="list-style-type: none"> ● Known inflammatory bowel disease and none of the above critical factors but flare of symptoms ● Suspected inflammatory bowel disease where chronic diarrhoea or other suspicious symptoms > 6 weeks with elevated faecal calprotectin (> 100 mcg/g), and none of the above critical factors or symptoms
<p>Category 3 (clinically recommended to be seen within 365 calendar days)</p>	<ul style="list-style-type: none"> ● Longstanding bowel symptoms suggestive of irritable bowel syndrome with none of the above critical factors or laboratory/imaging changes ● Lower gastrointestinal symptoms (e.g. suspected pancreatic exocrine insufficiency, suspected cholangitis colitis, faecal incontinence) and seeking specialist advice
<p>Referral requirements</p>	
<ul style="list-style-type: none"> ● Stool MC and S, oral contraceptive pill (OCP) and Clostridium difficile toxin (Category 1 referrals only) ● Molecular testing of stool for infection ● Personal or family history of inflammatory bowel disease ● Full blood count ● Faecal calprotectin result ● Liver function test result ● Electrolytes, urea and creatinine (EUC) ● Iron studies ● C-reactive protein (CRP) ● Relevant imaging reports ● Current and previous colonoscopy results ● Smoking history 	
<p>Other content</p>	
<ul style="list-style-type: none"> ● If the patient identifies as Aboriginal and/or Torres Strait Islander ● If the patient is considered 'at risk' and/or among a vulnerable, disadvantaged or priority population 	

Liver dysfunction (adult)

Emergency	
<p><i>If any of the following are present or suspected, please refer the patient to the emergency department (via ambulance if necessary) or seek emergent medical advice via phone to on-call consultant/registrar.</i></p> <ul style="list-style-type: none"> Acute liver failure (e.g. encephalopathy with liver test severe derangement, especially INR > 1.5 or bilirubin > 60) Severe hepatic encephalopathy 	
Out of scope / not routinely provided	
<ul style="list-style-type: none"> Fatty liver with normal liver function tests and Fibrosis-4 (FIB 4) score below 1.26 	
Access / prioritisation	
<p>Category 1 (clinically recommended to be seen within 30 calendar days)</p>	<ul style="list-style-type: none"> Abnormal liver function tests if any of the following: <ul style="list-style-type: none"> Bilirubin > 34 Albumin < 35 INR > 1.7 and/or platelets are outside normal range in setting of known or suspected liver disease (excluding unconjugated hyperbilirubinemia) Abnormal liver function tests associated with new symptoms such as nausea, anorexia or $\geq 5\%$ unexplained weight loss in past 1 month or $\geq 10\%$ unexplained weight loss in past 6 months Persisting liver inflammation with ALT > 200 for > 1 month New, abnormal liver function in a pregnant woman
<p>Category 2 (clinically recommended to be seen within 90 calendar days)</p>	<ul style="list-style-type: none"> Liver disease treatment required that is outside the referrer scope of practice (e.g. viral hepatitis, haemochromatosis, autoimmune liver disease, primary sclerosing cholangitis, primary biliary cholangitis, Wilson’s disease) Metabolic syndrome or alcohol dependence suspected and non-invasive serological algorithm (Fibrosis-4 (FIB 4)) suggestive of cirrhosis (FIB 4 score above 3.5; a threshold where cirrhosis is likely), or elastography or radiologic evidence of cirrhosis
<p>Category 3 (clinically recommended to be seen within 365 calendar days)</p>	<ul style="list-style-type: none"> Metabolic syndrome associated liver function tests derangement suspected (MAFLD) and fibrosis indeterminate (FIB 4 score 1.3 – 3.5) where significant fibrosis and even cirrhosis is possible Abnormal liver function tests and negative liver screen regardless of aetiology, severity or degree of work-up performed Fibrosis assessment requested (Fibro scan referral) when FIB 4 score above 1.3 (indeterminate), or known / risk factors for chronic liver disease requiring monitoring

	(e.g. methotrexate, metabolic associated steatohepatitis, Hepatitis B)
Referral requirements	
<ul style="list-style-type: none"> • Liver function test results (current and previous) • Full blood count/Coagulation profile • Upper abdomen ultrasound • Hepatitis A serology (HAV IgG) • Hepatitis B serology (HBV sAg, sAb, core IgG) • Hepatitis C serology (HCV Ab) • Iron studies • HbA1c • Medication list, including over the counter and herbal medicines • Alcohol intake (duration and quantity) • Risk factors for viral hepatitis 	
Other content	
<ul style="list-style-type: none"> • Previous ultrasound, CT or MRI reports • Elastography or other liver imaging • Vaccination history • Any relevant family history • Additional pathology tests (e.g. autoimmune hepatitis, primary biliary cirrhosis, Wilson’s disease, Genetics disorders) • If the patient identifies as Aboriginal and/or Torres Strait Islander • If the patient is considered ‘at risk’ and/or among a vulnerable, disadvantaged or priority population 	

Cirrhosis (suspected or known) (adult)

Emergency	
<p>If any of the following are present or suspected, please refer the patient to the emergency department (via ambulance if necessary) or seek emergent medical advice via phone to on-call consultant/registrar.</p> <ul style="list-style-type: none"> Suspected or known cirrhosis with acute clinical decompensation event (e.g. encephalopathy, gastrointestinal bleeding or new onset ascites, especially if with pain, fever or other systemic symptoms) Acute liver failure (e.g. encephalopathy with liver test severe derangement, especially INR > 1.5 or bilirubin > 60) Sepsis in a patient with cirrhosis Suspected ascending cholangitis Severe hepatic encephalopathy Suspected spontaneous bacterial peritonitis 	
Out of scope / not routinely provided	
<ul style="list-style-type: none"> Nil 	
Access / prioritisation	
<p>Category 1 (clinically recommended to be seen within 30 calendar days)</p>	<ul style="list-style-type: none"> Suspected or known cirrhosis on imaging, non-invasive testing or pathology, and any of the following are present: <ul style="list-style-type: none"> Decompensation event (current or previous) (e.g. history of jaundice, ascites, confusion, gastrointestinal bleeding) High risk features on laboratory (Child-Pugh > B7, Model for End-stage Liver Disease (MELD) > 9, platelets < 150) Elevated alpha-fetoprotein (AFP) or space occupying lesion on imaging
<p>Category 2 (clinically recommended to be seen within 90 calendar days)</p>	<ul style="list-style-type: none"> Suspected or known cirrhosis on imaging, non-invasive testing or bloods, and any of the following are present: <ul style="list-style-type: none"> Underlying liver disease identified which cannot be treated in primary care (e.g. viral hepatitis – discretion of referring clinician, haemochromatosis, autoimmune liver disease, primary sclerosing cholangitis) New diagnosis of cirrhosis with no prior specialist assessment or management plan
<p>Category 3 (clinically recommended to be seen within 365 calendar days)</p>	<ul style="list-style-type: none"> Routine care for known cirrhosis which requires assessment within 6 months (e.g. hepatocellular cancer surveillance, oesophageal varices screening, nutrition) <i>(patients with stable compensated who only require hepatocellular cancer surveillance may be appropriately managed in general practice)</i>

Referral requirements

- Alcohol intake
- Risk factors for viral hepatitis
- Medication list, including over the counter and herbal medicines
- Current and previous liver function tests
- Full blood count/Coagulation profile
- Upper abdomen ultrasound
- Hepatitis A serology (HAV IgG)
- Hepatitis B serology (HBV sAg, sAb, core IgG)
- Hepatitis C serology (HCV Ab)
- Iron studies

Other content

- HbA1c
- ANA, immunoglobins
- Previous ultrasound, CT or MRI reports
- Previous endoscopy reports
- Vaccination history
- Any relevant family history
- If the patient identifies as Aboriginal and/or Torres Strait Islander
- If the patient is considered 'at risk' and/or among a vulnerable, disadvantaged or priority population

Upper gastrointestinal dysfunction (adult)

Emergency	
<p><i>If any of the following are present or suspected, please refer the patient to the emergency department (via ambulance if necessary) or seek emergent medical advice via phone to on-call consultant/registrar.</i></p> <ul style="list-style-type: none"> Dysphagia with obstruction from food or foreign body associated with inability to tolerate oral intake, especially if drooling or pooling of saliva 	
Out of scope / not routinely provided	
<ul style="list-style-type: none"> Patients with controlled symptoms Non-smoker and < 50 years old that ceases treatment and symptoms return Belching Halitosis 	
Access / prioritisation	
<p>Category 1 (clinically recommended to be seen within 30 calendar days)</p>	<ul style="list-style-type: none"> Dysphagia any age or symptoms (excluding longstanding) Dyspepsia/heartburn/reflux with any of the following present: <ul style="list-style-type: none"> Warning symptoms: <ul style="list-style-type: none"> Haematemesis or melaena ≥ 5% unexplained weight loss in past 1 month or ≥ 10% unexplained weight loss in past 6 months Laboratory indicators: <ul style="list-style-type: none"> Iron deficiency Anaemia Low albumin Abnormal imaging of the upper gastrointestinal tract suggesting space occupying lesion
<p>Category 2 (clinically recommended to be seen within 90 calendar days)</p>	<ul style="list-style-type: none"> Helicobacter eradication failed (urea breath test) after standard 1st line therapy (7-14 days triple esomeprazole, clarithromycin, amoxicillin or metronidazole) > 50 years old with new, unexplained or persistent dyspepsia symptoms despite proton pump inhibitors
<p>Category 3 (clinically recommended to be seen within 365 calendar days)</p>	<ul style="list-style-type: none"> < 49 years old with longstanding, unexplained or persistent dyspepsia despite proton pump inhibitors Screening for Barrett’s oesophagus in patients with longstanding gastroesophageal reflux without additional symptoms Surveillance of known Barrett’s oesophagus Surveillance for gastric cancer in patients with history of intestinal metaplasia

Referral requirements

- Family history of gastrointestinal cancers
- History of weight loss
- Duration of symptoms
- Medical management to date (e.g. proton pump inhibitors, helicobacter pylori treatment)
- Full blood count
- Iron studies
- Smoking history

Other content

- Previous endoscopy/histopathology results
- Relevant imaging reports
- Helicobacter pylori results, including urea breath tests
- If the patient identifies as Aboriginal and/or Torres Strait Islander
- If the patient is considered 'at risk' and/or among a vulnerable, disadvantaged or priority population